

Selective Labeling Of Proteins On Their N-Termini With Synthetic Peptides

Tech ID: 23345 / UC Case 2006-124-0

BACKGROUND

Direct and selective labeling of α -amines or α -carboxylates of the N-terminal amino acid in proteins is a powerful approach for profiling proteins in complex mixtures. Approximately 80% of mammalian proteins are N-terminally acetylated. Thus, N-terminal labeling provides greater signal over background than does C-terminal labeling. However, such labeling must still be extremely selective for α -amines over lysine ϵ -amines, which are approximately 25 times more abundant in an average protein.

TECHNOLOGY DESCRIPTION

Investigators at UCSF have invented a general method to selectively label proteins on their N-termini with synthetic peptides. Briefly, this method uses an engineered enzyme subtiligase and a peptide substrate specially tailored to the proteomic workflow. The substrate comprises a peptide glycolate ester with a subtiligase cleavage site, a biotin label, and a TEV-cleavage site. The action of the subtiligase enzyme results in proteins that are selectively biotinylated on the N-terminus. The biotin label can then be used to capture these labeled proteins from a complex mixture using immobilized avidin. Finally, the TEV cleavage site allows for the enrichment of captured proteins or peptides for downstream analysis such as mass spectroscopy. Other N-terminal labels can also be used in place of biotin making the process very versatile.

APPLICATION

- Specific labeling of α -amines of N-terminal amino acid in proteins
- Global proteomic profiling of complex mixtures of proteins and polypeptides.

ADVANTAGES

- High specificity for the N-terminus of proteins
- Selective labeling of α -amines over lysine ϵ -amines of proteins

STAGE OF DEVELOPMENT

Pre-clinical

CONTACT

Kristin A. Agopian
kristin.agopian@ucsf.edu
tel: 415-340-2619.



OTHER INFORMATION

KEYWORDS

protein labeling, proteomics,
N-termini labeling, protein
profiling, synthetic peptides

CATEGORIZED AS

- [Biotechnology](#)
- [Proteomics](#)

RELATED CASES

2006-124-0, 2017-035-0

RELATED MATERIALS

- ▶ Mahrus S, et al. Global Sequencing of Proteolytic Cleavage Sites in Apoptosis by Specific Labeling of Protein N Termini. Cell 134, 866–876, September 5, 2008
- ▶ Wildes, D & Wells, J.A. (2010) Proc. Natl. Acad. Sci. USA 107(10), 4561-4566. “Sampling the N-terminal proteome of human blood”. PMID: 20173099
- ▶ Agard, N.J, et al (2010) Mol. Cell. Proteomics 5, 880-893. “Inflammatory stimuli regulate caspase substrate profiles”. PMID: 20173201
- ▶ Agard, N.J., et al (2012) Proc Natl Acad Sci U S A. 109:1913-8. “Global kinetic analysis of proteolysis via quantitative targeted proteomics”. PMID: 22308409
- ▶ Crawford, E.D., Seaman, J.E., et al (2012), “The DegraBase: A database of proteolysis in healthy and apoptotic human cells”, Mol Cell Proteomics

PATENT STATUS

Country	Type	Number	Dated	Case
United States Of America	Issued Patent	8,679,771	03/25/2014	2006-124

ADDRESS

UCSF
Innovation Ventures
600 16th St, Genentech Hall, S-272,
San Francisco,CA 94158

CONTACT

Tel:
innovation@ucsf.edu
https://innovation.ucsf.edu
Fax:

CONNECT

 Follow  Connect

© 2013 - 2022, The Regents of the University
of California
[Terms of use](#) [Privacy Notice](#)