

Microbial-Induced Barriers To Striga Parasitism

Tech ID: 33538 / UC Case 2023-530-0

ABSTRACT

Researchers at the University of California, Davis have discovered an *Arthrobacter* bacterial strain that promotes suberization of the endodermis in sorghum roots. Suberin, a poly-fatty acid polymer, acts as a physical barrier in sorghum roots, helping to prevent infection by the parasitic plant *Striga hermonthica*, a significant threat to sorghum production. These microbial-based solutions offer a cost-effective and easily deployable strategy to manage *Striga* infection in the predominantly smallholder farmer-driven sorghum cultivation of sub-Saharan Africa.

FULL DESCRIPTION

Sorghum, a vital cereal crop in sub-Saharan Africa, frequently faces significant yield losses due to infestations by the parasitic weed *Striga hermonthica*. *Striga* attaches to sorghum roots and penetrates their tissues, causing reduced water and nutrient uptake, leading to wilting, stunted growth, and decreased yields. Existing *Striga* management methods have proven inefficient and challenging to integrate into traditional agricultural practices.

This technology identifies a novel approach to combat *Striga* infection by harnessing the power of beneficial soil microbes. UC Davis researchers discovered a soil microbiome that suppressed *Striga* infection in sorghum by increasing suberin content in the plant's root system. Mathematical modeling predicts the specific microbial genera responsible for this suppression. Among the selected candidates, an *Arthrobacter* strains was highly effective at inducing suberization of endodermis and exodermis, which strengthened the suberin barrier in sorghum roots.

APPLICATIONS

- ▶ **Sorghum Agriculture:** This technology can be applied to sorghum cultivation in sub-Saharan Africa, providing a cost-effective and accessible method for managing *Striga hermonthica* infestations, thus improving crop yields and food security.
- ▶ **Microbial Inoculants:** The *Arthrobacter* strain can be developed into microbial inoculants applied to sorghum seeds, seedlings, or soil to enhance the suberin barrier and protect the crop from *Striga* infestation.
- ▶ **Plant Biotechnology:** Advances the understanding of suberin deposition in plant roots, potentially leading to the development of genetically enhanced sorghum varieties with stronger suberin barriers.

FEATURES/BENEFITS

- ▶ **Effective Striga Management:** The technology offers an effective and sustainable solution for controlling *Striga hermonthica* infestations in sorghum, reducing crop losses, and improving food production.

CONTACT

Eugene Sisman

esisman@ucdavis.edu

tel: 530-754-7650.



INVENTORS

- ▶ Brady, Siobhan
- ▶ Kawa, Dorota

OTHER INFORMATION

KEYWORDS

sorghum, striga,
microbial inoculants, soil
microbiome, crop
protection, plant
biotechnology

CATEGORIZED AS

- ▶ **Agriculture & Animal Science**
 - ▶ Plant Traits
- ▶ **Biotechnology**
 - ▶ Food
 - ▶ Other

RELATED CASES

2023-530-0

► Affordable and User-Friendly: Microbial-based solutions are cost-effective and easy to implement, making them particularly suitable for smallholder farmers in sub-Saharan Africa.

PATENT STATUS

Country	Type	Number	Dated	Case
Patent Cooperation Treaty	Published Application	WO 2024/092266	05/02/2024	2023-530

Additional Patent Pending

ADDITIONAL TECHNOLOGIES BY THESE INVENTORS

► [Genes Controlling Barrier Formation in Roots](#)

University of California, Davis

Technology Transfer Office

1 Shields Avenue, Mrak Hall 4th Floor,
Davis,CA 95616

Tel:
530.754.8649
techtransfer@ucdavis.edu
<https://research.ucdavis.edu/technology-transfer/>
Fax:
530.754.7620

© 2024, The Regents of the University of California
[Terms of use](#)
[Privacy Notice](#)