Request Information Permalink

MONOCLONAL ANTIBODIES SPECIFIC FOR TRIAZINE HERBICIDES INCLUDING ATRAZINE.

Tech ID: 18144 / UC Case 1990-004-0

ABSTRACT

Simazine and atrazine haptens with mercaptopropionic acid and aminohexanoic acid spacers were synthesized and conjugated to proteins via N-hydroxysuccinimide active esters. Mabs derived from immunized mice had I50 values of 3 to 4 ppb. The EIAs are compatible with simplified methods for triazine extraction and concentration from soil and water. The limit of detection for atrazine was 0.05 to 0.1 ppb.

Reference;

Karu et al. Monoclonal Immunoassay of Triazine Herbicides Development and Implementation 1991. ACS Symposium Series No. 451 Immunochemical Methods for Trace Chemical Analysis. pp59-77.

CONTACT

Javed Afzal jafzal@berkeley.edu tel: 510-643-7201.



OTHER INFORMATION

KEYWORDS

research tool, antibody

CATEGORIZED AS

» Research Tools

» Antibodies

RELATED CASES1990-004-0



University of California, Berkeley Office of Technology Licensing

2150 Shattuck Avenue, Suite 510, Berkeley,CA 94704

Tel: 510.643.7201 | Fax: 510.642.4566

 $ipira.berkeley.edu/\mid otl\mbox{-feedback@lists.berkeley.edu}$

© 2009 - 2010, The Regents of the University of California

Terms of use | Privacy Notice