# TRANSCRIPTIONAL ACTIVATION FACTORS (CJUN, AP-2, SP1, CTF/NF-1)

Tech ID: 18327 / UC Case 1994-062-0

## ABSTRACT

Eukaryotic promoters are regulated by a combination of sequence-specific DNA-binding proteins, general transcription initiation factors, and associated accessory factors. The sequence-specific transcription factors can be divided into several classes on the basis of the activation domains they posess. This disclosure relates to several Human cDNA clones and/or expression vectors encoding c-jun, AP2, SP1, and CTF/NF1 TAF genes.

Reference;

B.D. Dynlacht, et al., Isolation of Coactivators Associated with the TATA-Binding Protein That Mediate Transcriptional Activation. 1991. Cell 66:563-76

## CONTACT

Craig K. Kennedy
craig.kennedy@berkeley.edu
tel: .

INTRODUCING
UC TechAlerts
New technology matches delivered to your email at your preferred schedule
SEARCH SAVE SEARCH
Learn More

## OTHER INFORMATION

**KEYWORDS**

research tool, expression system, vector

**CATEGORIZED AS**

» **Research Tools**

» Expression System

» Vectors

**RELATED CASES**

1994-062-0

## RELATED TECHNOLOGIES

▶ Hybridomas Producing Monoclonal Antibodies To Basal Transcription Factors


© 2009 - 2010, The Regents of the University of California
Terms of use | Privacy Notice