

[Request Information](#)

[Permalink](#)

## Phasor method to fluorescence lifetime microscopy discriminates metabolic states of germ cells in a living tissue / Phase method to fluorescence lifetime microscopy to discriminate metabolic state of cells in living tissue

Tech ID: 21811 / UC Case 2010-805-0

### BRIEF DESCRIPTION

Researchers at UC Irvine have developed a label-free way to discriminate the *in vivo* metabolic state of cells in a tissue through phasor analysis. This provides a sensitive, non-invasive optical tool to study cell differentiation and disease progression, predict cell fate, and for use in cell sorting applications of unlabeled tissues.

### FULL DESCRIPTION

It is of interest to study cell metabolic states. Autofluorescence in live tissues, which arise from endogenous proteins and physiologically relevant fluorophores such as collagen, elastin, porphyrin, retinoids, flavins, nicotinamide adenine dinucleotide, hemoglobin, and serotonin, can give indication of cell metabolic activity. Furthermore, NADH and FAD are the main metabolic coenzymes involved in oxidative phosphorylation and glycolysis, reporting on metabolic changes associated with cell carcinogenesis and differentiation. Non-invasive optical methods, such as multi-photon microscopy, have been developed for high-resolution and long-term imaging of living tissues. However, the ability to assign autofluorescence to specific tissue components or molecular sources using standard multi-photon microscopy cannot be done. Other methods to assign autofluorescence to specific tissue components, including principle component analysis and multi-exponential fitting, fail to capture the complexity of living tissues.

Researchers at UC Irvine have developed a sensitive, label-free way to discriminate *in vivo* metabolic states of cells in a tissue using the phasor approach to fluorescence lifetime imaging microscopy (FLIM). This phasor approach to FLIM relies on separating and identifying tissue components by cluster analysis.

### CONTACT

Alvin Viray  
aviray@uci.edu  
tel: 949-824-3104.



### OTHER INFORMATION

#### CATEGORIZED AS

- » **Biotechnology**
- » Other
- » **Imaging**
- » Molecular
- » Other
- » **Medical**
- » Imaging
- » Research Tools
- » Screening
- » **Research Tools**
- » Other

### RELATED CASES

2010-805-0




Figure 1.

Figure 1 illustrates an example of phasor analysis of FLIM tissue images to identify tissue components. Figure 1a shows the phasor locations of pure chemical species for green fluorescent protein (GFP) in Tris buffer, Retinol in DMSO (pH 8.5), Retinoic acid in DMSO (pH 8.5), FAD in water (pH 7.4), free NADH in Mops buffer (pH 7), bound NADH in Mops buffer (pH 7) and lactate dehydrogenase, Protoporphyrin IX in dimethylformamide:methanol (pH 7). Figure 1b is the fluorescent intensity image of a seminiferous tubule from a mouse expressing GFP from an Oct4 transgene. Figure 1c is the phasor plot of the FLIM image acquired in Figure 1b. Figure 1d is the phasor color map, where the color in the phasor color map correspond to the color of the cluster of the phasor plot shown in Figure 1c.




Figure 2.

The inventors have demonstrated the ability to identify the metabolic states of germ cells during differentiation using the phasor approach to FLIM. Figure 2a shows metabolite gradients in *C. Elegans* Germ line when excited at 740 nm. Figure 2b shows phasor plot selection using linear cluster that represents all possible relative concentrations of pure FAD (red), free NADH (blue), and NADH bound to MDH (green). Figure 2c shows the spectral image of the *C. Elegans* excited at 740 nm in the same field of view with R1 and R2 indicating regions of interest. The emission spectra R1 and R2 is shown in Figure 2d. The blue shift of the spectrum indicates an increase in bound NADH with respect to free NADH during differentiation.

This method allows for straightforward interpretation of intrinsic fluorescence signal from living tissues. Images of molecule species are obtained through their finger prints without the need to resolve and assign exponential components to the fluorescence species.

## SUGGESTED USES

This invention can be used to monitor metabolic states of cells, in particular stem cells, both in vitro and in vivo, and study cell fates. This invention can also be used to differentiate tissue composition.

## ADVANTAGES

The invention uses the phasor approach for FLIM. This is a label-free method that allows for identification and quantification of relative concentrations of molecular species in living tissues.

## PATENT STATUS

Country	Type	Number	Dated	Case
United States Of America	Issued Patent	10,222,335	03/05/2019	2010-805

**UCI** Beall  
Applied Innovation

5270 California Avenue / Irvine, CA  
92697-7700 / Tel: 949.824.2683



© 2011 - 2019, The Regents of the University of  
California  
[Terms of use](#)  
[Privacy Notice](#)