Request Information

INTRINSIC NAVIGATION FROM VELOCITY-ENCODING GRADIENTS IN PHASE-CONTRAST MRI

Tech ID: 24756 / UC Case 2015-097-0

PATENT STATUS

Country	Туре	Number	Dated	Case
United States Of America	Issued Patent	10,132,902	10/20/2018	2015-097

BRIEF DESCRIPTION

Time-resolved phase contrast MRI (4D flow) can quantify cardiac function and flow. The technique may even permit complex anatomical assessment, thus comprising a comprehensive exam in a single scan. Unfortunately, artifacts from respiratory motion compromise this ability. Therefore, we developed a simple method to measure motion using readily available navigation information from the velocity-encoding gradients without any significant modification to conventional sequences.

SUGGESTED USES

>> Can be used in conjunction with any type of phase-contrast MR imaging scheme and any type of reconstruction strategy.

» Allows for high-quality images to be reconstructed from exams with long acquisition times and other types of exams that are sensitive to motion defects.

» Can be used for any part of the body that requires assessment using phase-contrast imaging. It is especially useful in cases when patients have

difficulty holding still for long time periods.

>>> For longer scans (>1 min), breath-holds are no longer possible and respiratory motion must be considered. With the navigation technique, this imaging

modality can be made accessible to a wider patient population.

ADVANTAGES

- » In conventional phase-contrast sequences, no alterations to the sequence timing or gradient waveform are needed.
- » Navigator information is available for every repetition time and is naturally synchronized to the acquired imaging data.
- » In conjunction with a with a high-density multi-channel coil receiver, each coil element provides spatial localization to the navigator signals.
- » The navigator measured can be used for motion compensation purposes or to prospectively gate the acquisition.
- >>> The method can be used for any sampling strategy (both Cartesian and non-Cartesian).

RELATED MATERIALS

» Intrinsic Navigation from Velocity-Encoding Gradients in Phase-Contrast MRI - 12/07/2015

ADDITIONAL TECHNOLOGIES BY THESE INVENTORS

Method and Device for Designing Smooth Sequences of Spoke Endpoints in MRI

Method for Motion Sensing in MRI Using Preamplifier RF Intermodulation

CONTACT

Terri Sale terri.sale@berkeley.edu tel: 510-643-4219.



INVENTORS

» Lustig, Shimon M.

OTHER INFORMATION

KEYWORDS

healthcare, imaging, MRI

CATEGORIZED AS

» Imaging

» Medical

» Medical

>> Imaging

RELATED CASES 2015-097-0

Permalink



University of California, Berkeley Office of Technology Licensing 2150 Shattuck Avenue, Suite 510, Berkeley,CA 94704 Tel: 510.643.7201 | Fax: 510.642.4566 https://ipira.berkeley.edu/ | otl-feedback@lists.berkeley.edu © 2015 - 2018, The Regents of the University of California Terms of use | Privacy Notice