

Request Information

Permalink

Monoclonal Antibody Against PNPase (Clone 3H5)

Tech ID: 25121 / UC Case 2011-375-0

BRIEF DESCRIPTION

Mouse monoclonal antibody against the human mitochondrial polyribonucleotide nucleotidyltransferase 1 (PNPase). This antibody has been tested for use in immunocytochemistry/immunofluorescence, immunoprecipitation, and western blot.

FULL DESCRIPTION

RNA-binding protein implicated in numerous RNA metabolic processes. Catalyzes the phosphorolysis of single-stranded polyribonucleotides processively in the 3'-to-5' direction. Mitochondrial intermembrane factor with RNA-processing exoribonuclease activity. Component of the mitochondrial degradosome (mtEXO) complex, that degrades 3' overhang double-stranded RNA with a 3'-to-5' directionality in an ATP-dependent manner. Required for correct processing and polyadenylation of mitochondrial mRNAs. Plays a role as a cytoplasmic RNA import factor that mediates the translocation of small RNA components, like the 5S RNA, the RNA subunit of ribonuclease P and the mitochondrial RNA-processing (MRP) RNA, into the mitochondrial matrix. Plays a role in mitochondrial morphogenesis and respiration; regulates the expression of the electron transport chain (ETC) components at the mRNA and protein levels. In the cytoplasm, shows a 3'-to-5' exoribonuclease mediating mRNA degradation activity; degrades c-myc mRNA upon treatment with IFN β /IFN-beta, resulting in a growth arrest in melanoma cells. Regulates the stability of specific mature miRNAs in melanoma cells; specifically and selectively degrades miR-221, preferentially. Plays also a role in RNA cell surveillance by cleaning up oxidized RNAs. Binds to the RNA subunit of ribonuclease P, MRP RNA and miR-221 microRNA.

The described mouse monoclonal antibody binds to human mtPAP and has been described for use in immunocytochemistry/immunofluorescence, immunoprecipitation, and western blot.

SUGGESTED USES

Immunocytochemistry/immunofluorescence (1:100-1:1000)* Immunoprecipitation Western blot

* Suggested dilutions

ADVANTAGES

Specific antibody to detect mtPAP (clone 1D3) by antibody-antigen complex formation.

RELATED MATERIALS

» Protein-protein interactions between human exosome components support the assembly of RNase PH-type subunits into a six-membered PNPase-like ring. - 11/01/2002

CONTACT

Patricia H. Chan
patricia.chan@uci.edu
tel: 949-824-6821.



INVENTORS

- » Guo, Xuning
- » Lee, Wen Hwa

OTHER INFORMATION

KEYWORDS

Monoclonal Antibody,
Centrosomal protein 164kDa,
Cep 164,
Immunocytochemistry/immunofluorescence,
Immunoprecipitation,
Western blot

CATEGORIZED AS

- » **Agriculture & Animal Science**
 - » Animal Science
- » **Biotechnology**
 - » Genomics
 - » Health
 - » Proteomics

» **Imaging**

- » Medical
- » Molecular

» **Materials & Chemicals**

- » Biological

» **Medical**

- » Diagnostics
- » Disease:
Autoimmune and
Inflammation
- » Disease: Blood and
Lymphatic System
- » Disease: Cancer
- » Disease:
Cardiovascular and
Circulatory System
- » Disease: Central
Nervous System
- » Research Tools
- » Therapeutics
- » Vaccines

» **Nanotechnology**

- » Tools and Devices

» **Research Tools**

- » Antibodies

» **Sensors & Instrumentation**

- » Medical
- » Scientific/Research

RELATED CASES

2011-375-0, 2011-387-0,
2011-386-0, 2011-385-0,
2011-384-0, 2011-383-0,
2011-382-0

ADDITIONAL TECHNOLOGIES BY THESE INVENTORS

- ▶ Monoclonal Antibody Against PNPase (Clone 4C11)
- ▶ Monoclonal Antibody Against mtPAP (Clone 3D2)

UCI Beall
Applied Innovation

5270 California Avenue / Irvine, CA
92697-7700 / Tel: 949.824.2683



© 2015, The Regents of the University of
California
Terms of use
Privacy Notice