Find Technologies

Permalink

Request Information

Substantial defluorination of chlorinated PFCAs

Tech ID: 33655 / UC Case 2024-706-0

FULL DESCRIPTION

Background

Polychlorofluoroalkyl compounds represent a large group of per- and polyfluoroalkyl substances (PFAS). Well known examples include polychlorotrifluoroethylene (PCTFE) and chlorotrifluoroethylene (CTFE). The thermolysis of PCTFE and the metabolism of CTFE results in the formation of CTFE oligomer carboxylic acids which pose a risk to the hydrosphere. Prior research examining the biodegradation of chlorinated polychlorofluoroalkyl substances (CI-PFAS) focused on a narrow range of compounds and identified reductive chlorination as the primary mechanism. There remains a significant gap regarding the biodegradability and ultimate fate of CI-PFAS in the environment.

Technology

Prof. Yujie Men and her team investigated the biotransformation of CI-PFAS by anaerobic microbes and have recently demonstrated that microbial dechlorination specifically hydrolytic dechlorination is a crucial step in the defluorination of CTFE oligomers. They also demonstrate that increasing the number of chlorine substitutions enhances their biodegradability leading to more extensive defluorination. The team shows that an anaerobic-aerobic treatment process could enhance the complete destruction of CI-PFAS.



CONTACT

Venkata S. Krishnamurty venkata.krishnamurty@ucr.edu tel: .

OTHER INFORMATION

KEYWORDS anaerobic, biodegradation, CTFE, Cl-PFCA, hydrolytic dechlorination, microbial dechlorination, PFAS, wastewater treatment

CATEGORIZED AS

Biotechnology
Other

- Environment
 - Remediation

RELATED CASES 2024-706-0

Image showing the biotransformation pathways. Green arrows represent dechlorinating reactions. Red arrows represent defluorinating

reactions. The thickness of the red arrows represents the number of fluoride ions (F) released - a metric for the extent of breakdown of carbon-

fluorine (C-F) bonds.



Graph showing the parent compound decay and F⁻ formation. The data are represented as mean values. The error bars represent the standard deviation from n=3 replicate experiments.

ADVANTAGES

Cost-effective bioremediation - Sequential anaerobic-aerobic bioreactors, commonly used in wastewater treatment, could be adapted for a cost-effective cleanup of impacted environments.

Designing safer alternatives - Incorporating chlorine substitutions in PFAS molecules could offer a viable strategy for designing new PFAS substances with similar functions but a significantly reduced environmental footprint.

Understanding environmental fate - the technology can be used to predict the behavior and persistence of CI-PFAS in anaerobic environments.

Enhanced PFAS source tracking - the technology developed can aid in tracking the origin and fate of PFAS in the environment.

APPLICATIONS

- Cost-effective remediation of PFAS contamination.
- Design of safer PFAS alternatives.

INVENTOR INFORMATION

Please review all inventions by Prof. Men and her team at UCR.

Please visit Prof. Men's group website to learn more about their research.

RELATED MATERIALS

Substantial defluorination of polychlorofluorocarboxylic acids triggered by anaerobic microbial hydrolytic dechlorination

PATENT STATUS

Patent Pending

University of California, Riverside Office of Technology Commercialization 200 University Office Building, Riverside,CA 92521 otc@ucr.edu https://research.ucr.edu/

Terms of use | Privacy Notice | © 2024, The Regents of the University of California