

## B-raf/loxP-flanked Mutant Mouse

Tech ID: 20279 / UC Case 2006-215-0

### BACKGROUND

B-raf is a member of the Raf family of intracellular signaling proteins, which also include Raf-1 and A-raf. Members of this family are involved in the all-important cellular signaling mechanism known as the Ras/Raf/MEK/ERK/MAPK signaling pathway. This ubiquitous pathway relays signals from outside the cell into the cells nucleus, regulating activities such as gene expression, differentiation, cell division, cell survival and cell death. B-raf regulates vital functions in the brain, testes, skin and bone marrow, and mutations in B-raf are found in malignant melanoma, certain types of thyroid and ovarian cancers, and sporadic types of colon cancers. Up to 6% of all human malignancies may harbor this mutation. Most recently, Sorafenib (Nexavar), which inhibits raf as well as a whole host of other signaling proteins, has been approved by the FDA for the treatment of advanced cancer of the kidney (clear-cell renal cell carcinoma).

### INNOVATION

UCLA researchers have made a B-raf/loxP-flanked transgenic mouse. Flanking the B-raf gene with the loxP sequence allows targeted excision of the B-raf gene only when the cell expresses the protein Cre. This allows selective deletion of B-raf in a cell or tissue-specific manner and in a time or developmentally-specific manner as well.

### APPLICATIONS

- ▶ Cancer research involving the Ras/Raf/MEK/ERK/MAPK signaling pathways
- ▶ Research into learning and memory, and hematopoiesis and myelopoiesis

### ADVANTAGES

- ▶ Uses proven Cre/LoxP DNA recombination technology for targeted gene excision
- ▶ Unlike knockout mutants with global B-raf deficiency, B-raf/loxP sequence allows deletion of the B-raf gene in a cell or tissue-specific manner
- ▶ Unlike knockout mutants, B-raf/loxP sequence allows deletion of the B-raf gene in a time or developmentally-specific manner, such as after tumorigenesis or transformation
- ▶ Allows functional testing in specific cell types without the confounding effects of deleting B-raf in unknown or unrelated cells and tissues

### RELATED MATERIALS

- ▶ Forebrain-specific knockout of B-raf kinase leads to deficits in hippocampal long-term potentiation, learning, and memory. *J Neurosci Res.* 2006

### ADDITIONAL TECHNOLOGIES BY THESE INVENTORS

- ▶ [Inducible Dominant Negative Disc1 Transgenic Mice as a Model for Schizophrenia](#)
- ▶ [Phospho-specific Antibody for Cam Kinase II](#)
- ▶ [Statins as Treatment for Cognitive Dysfunction Associated with RASopathies](#)

### CONTACT

UCLA Technology Development Group  
ncd@tdg.ucla.edu  
tel: 310.794.0558.



### INVENTORS

- ▶ Silva, Alcino J.

### OTHER INFORMATION

#### KEYWORDS

research tools therapeutics, mouse model, mice, ERK, cancer, conditional, hematopoiesis, myelopoiesis, learning and memory

#### CATEGORIZED AS

- ▶ [Research Tools](#)
- ▶ [Animal Models](#)

#### RELATED CASES

2006-215-0

**UCLA Technology Development Group**

10889 Wilshire Blvd., Suite 920, Los Angeles, CA 90095

[tdg.ucla.edu](http://tdg.ucla.edu)

Tel: 310.794.0558 | Fax: 310.794.0638 | [ncd@tdg.ucla.edu](mailto:ncd@tdg.ucla.edu)

© 2013 - 2014, The Regents of the University of California

[Terms of use](#)

[Privacy Notice](#)

